

Comment on the Background Independent Open String Theory

Samson L. Shatashvili [†] [#]

School of Natural Sciences
Institute for Advanced Study
Olden Lane
Princeton, NJ 08540

A direct derivation of the string field theory action in the Witten's background independent open string theory for the case when ghosts and matter are decoupled is given and consequences are discussed.

March, 92

[†] Research supported by DOE grant DE-FG02-90ER40542.

[#] On leave of absence from St. Petersburg Branch of Mathematical Institute (LOMI), Fontanka 27, St.Petersburg 191011, Russia.

Recently, a background independent formulation of open string field theory was given in [1]. This theory is formally defined in the space of all two dimensional world sheet theories, with the world sheet lagrangian given by

$$L = \int_{\Sigma} d^2z \left(\frac{1}{8\pi} g^{\alpha\beta} \partial_{\alpha} X^i \partial_{\beta} X^j \eta_{ij} + \frac{1}{2\pi} b^{ij} D_i c_j \right) + \int_{\partial\Sigma} d\theta V(X, b, c) \quad (1)$$

Here, the first term is the closed string background and the second term describes an arbitrary boundary interaction with the condition that the boundary operator V has the form¹

$$V = b_{-1} O, \quad (2)$$

where O is a general operator of the ghost number 1. If we denote by O_i some basis in the space of such operators, $O = \omega^i O_i$, the basic definition of the string field theory action S can be given via an equation that involves the two-point function of O_i 's on the world-sheet:

$$dS = \frac{1}{2} \int_0^{2\pi} d\theta_1 d\theta_2 < dO(\theta_1) \{Q, O\}(\theta_2) >, \quad (3)$$

where $dS = d\omega^i d/dt^i S$ and $dO = d\omega^i O_i$ and $< \dots >$ denotes unnormalized correlation function.²

In this letter we will show that if the operator V doesn't contain ghosts and is a function of purely bosonic variables, the right hand side of (3) actually reduces to a one-point function $d < \Omega^i V_i >$ with Ω being a certain linear combination of couplings ω . From this statement it follows immediately that the left hand side of (3) is indeed an exact form and that

$$S = (\Omega^i \frac{d}{d\omega^i} + 1) Z, \quad (4)$$

where Z is the world sheet partition function. We will give the expression for Ω below (see (15), (16), (17)).

The fact that the action can be written in the form (4) was demonstrated before in [2] by explicit calculations for the case of quadratic interaction $V = \omega X^2$; also in [2] for

¹ For definition of b_{-1} see [1].

² It can be shown, using Ward identities, that the two point function on the right hand side in (3) is a closed form, see [1].

general $V(X)$ the existence of a relation of type (4) was proven and the algorithm for constructing the vector field in (4) was described.

Our approach differs from the one in [2]: we will not use the exactness of the one form on the right hand side of (3) but we will just reduce the correlation function in (3) to a total derivative of a one-point function. Also, we will give an explicit formula for the vector field in (4) and it's relation to the Virasoro generators on world sheet. As a consequence of the definition (3) we will see that the zeros of the vector field Ω , which define the string field theory equations of motion (see [2])³, are linear functions of couplings; therefore the equations of motion in the present formalism are also linear. At the end we will make some comments. The alternative derivation of the action functional presented here and the above observation might be helpful for a more detailed understanding of open string field theory.

To evaluate the correlation function in (3) we will write the BRST operator in terms of the matter and ghost stress tensors T_m, T_g :

$$Q = \int_0^{2\pi} d\theta c(\theta) [T_m(\theta) + T_g(\theta)]. \quad (5)$$

Also from the assumption that ghosts and matter are decoupled in the theory we can write $O = cV(X)$ and thus the correlation function is given by two terms

$$\begin{aligned} \frac{1}{2} \int_0^{2\pi} d\theta_1 d\theta_3 < c(\theta_1) dV(\theta_1) c(\theta_3) \partial_{\theta_3} c(\theta_3) V(\theta_3) > \\ + \frac{1}{2} \int_0^{2\pi} d\theta_1 d\theta_2 d\theta_3 < c(\theta_1) dV(\theta_1) c(\theta_2) c(\theta_3) [T_m(\theta_2), V(\theta_3)] > . \end{aligned} \quad (6)$$

To proceed further we need to evaluate the ghost correlation functions in (6). For the general 3-point function we have:

$$< c(\theta_1) c(\theta_2) c(\theta_3) > = 2(\sin(\theta_1 - \theta_2) - \sin(\theta_1 - \theta_3) + \sin(\theta_2 - \theta_3)). \quad (7)$$

This leads to a simple expression for the first correlator in (6)

³ The fact that the string field theory action on the classical equations of motion is given by the world-sheet bosonic partition function previously was suspected in [3] [4] and elaborated in terms of theory under the discussion in [2].

$$\frac{1}{2} \int_0^{2\pi} < d\theta_1 d\theta_2 (2 \cos(\theta_1 - \theta_2) - 2) < dV(\theta_1) V(\theta_2) >, \quad (8)$$

and to three terms for the second correlator in (6):

$$\begin{aligned} \frac{1}{2} \int_0^{2\pi} d\theta_1 d\theta_2 [i(- < e^{i\theta_1} dV(\theta_1) [L_{-1}, V(\theta_2)] > -c.c.) + \\ + i(- < dV(\theta_1) [L_1, e^{-i\theta_2} V(\theta_2)] > -c.c.) - \\ - 2 \sin(\theta_1 - \theta_2) < dV(\theta_1) [L_0, V(\theta_2)] >]. \end{aligned} \quad (9)$$

Here we use the notation $L_n = \int_0^{2\pi} d\theta e^{in\theta} T_m(\theta)$.

The last term in (9) can be simplified if we use the fact that $[L_0, V(\theta)] = \partial V(\theta)$ and thus after integration by parts over θ_2 it leads to

$$- \int_0^{2\pi} d\theta_1 d\theta_2 \cos(\theta_1 - \theta_2) < dV(\theta_1) V(\theta_2) >. \quad (10)$$

At the same time we can also simplify the first term in (9) just by using the observation that the commutator term here is nothing but the commutator of the Virasoro generator L_{-1} with the world sheet action (note that the bulk lagrangian commutes with all Virasoro generators). Thus in the path integral for this correlation function we could integrate by parts and because there is no anomaly in the measure (the construction of a background independent string field theory assumes that we have a critical string) we get:

$$\frac{i}{2} (< [L_{-1}, \int_0^{2\pi} e^{i\theta} dV(\theta)] > -c.c.). \quad (11)$$

Now we can combine (8), (9), (10) and (11) and due to some nice cancellations and the identity

$$< \int_0^{2\pi} d\theta_1 dV(\theta_1) \int_0^{2\pi} d\theta_2 V(\theta_2) > = d < \int_0^{2\pi} d\theta V(\theta) > - dZ, \quad (12)$$

we obtain

$$\begin{aligned} dZ - d < \int_0^{2\pi} d\theta V(\theta) > + \frac{i}{2} (< [L_{-1}, \int_0^{2\pi} d\theta e^{i\theta} dV(\theta)] > \\ - < \int_0^{2\pi} d\theta_1 dV(\theta_1) [L_1, \int_0^{2\pi} d\theta_2 e^{-i\theta_2} V(\theta_2)] > -c.c.), \end{aligned} \quad (13)$$

where Z denotes the world sheet partition function.

At this point we have to make an important assumption that the exterior derivative in the space of couplings, d , commutes with Virasoro generators $L_1, L_{-1}; [d, L] = 0$. This property was crucial also in the consideration of [1] (e.g. for the prove that the symplectic form is closed and BRST invariant, see [1]). It is important to keep in mind that, because we are dealing in principle with non-renormalizable theory and are using a cut-off, we have to require that short distance behavior of the exact Green function, $G(z_1, z_2)$, in the theory is still logarithmic; this is certainly true on the bulk, but not guaranteed when both z_1 and z_2 are on the boundary: $z_1 = \exp(i\theta_1), z_2 = \exp(i\theta_2)$. The latter requirement leads to certain restrictions on the couplings ω^i . Only in this case we can define the Virasoro algebra through the world sheet stress-tensor both in the bulk and on the boundary, and have geometric transformation properties for the boundary operators, corresponding to reparametrization of the world-sheet. Unfortunately, because generically we will have unrenormalizable boundary interactions, we can not make this point more elaborate. Although, we can make some comments if boundary interaction is quadratic:⁴ $V = \sum_0^\infty \omega_n^{ij} V_n^{ij} + a; V_n^{ij} = \int_0^{2\pi} d\theta X^i \partial^n X^j$, and a is a constant. In this case, Green function with both points on boundary is given by $\langle X^i(\theta_1) X^j(\theta_2) \rangle = \sum_{k=-\infty}^\infty (|k| + \sum_{n=0}^\infty \omega_n (ik)^n)_{ij}^{-1} \exp i(\theta_1 - \theta_2)$. Now if the infinite sum in the denominator for this Green function goes to a constant for large k , we will have a logarithmic short distance divergence and we can define all operators V by point splitting and subtracting the divergent part, which will not depend on the couplings.⁵

After the above comment we can now exchange the order of d and $L_{-1}(L_1)$ in (13), and thus conclude that the right hand side of (3) is an exact form and defines the string field theory action up to a constant:

$$d(Z - \langle \int_0^{2\pi} d\theta V(\theta) \rangle + \frac{i}{2} (\langle \int_0^{2\pi} d\theta e^{i\theta} [L_{-1}, V(\theta)] \rangle - c.c.)) = dS. \quad (14)$$

Thus, we have expressed the action S in terms of a one-point function of the boundary operator V . Let us represent V as a sum of basic operators, V_i , $V = \omega^i V_i$, where V_i corresponds to O_i introduced above and is a function of the field X and it's derivatives. The operation δ

⁴ I would like to thank E. Witten for pointing out this possibility.

⁵ The treatment of regularization problems in the case when there are finite number of non-zero couplings ω_n in quadratic potential is given in [5]

$$\delta V_i = \frac{i}{2}[e^{i\theta} L_{-1} - e^{-i\theta} L_1, V_i] \quad (15)$$

defines a linear transformation in the space of operators V_i , or equivalently, in the parameter space ω^i :

$$\delta(\omega^i V_i) = \hat{\omega}^i V_i = \omega^i \delta V_i, \quad (16)$$

and $\hat{\omega}$ is given by (15) and (16). This completes our derivation of desired formula for action S:

$$\begin{aligned} S &= (\Omega^i \frac{d}{d\omega^i} + 1)Z; \\ \Omega^i &= (\hat{\omega}^i - \omega^i), \end{aligned} \quad (17)$$

and zeros of Ω

$$\Omega^i = 0 \quad (18)$$

are classical equations of motion.

After the equations of motions (18) are derived for general boundary interactions (with restrictions on short distance behavior emphasized above) we would like to make some comments. As it follows from (18), (15) and (16), these equations are linear and gauge symmetries are given by zero modes of the operator $\delta - 1$. In the notation $V(X) = T(X) + A_\mu \partial X^\mu + \dots$ from (15) and (16) the equation $\Omega = 0$ leads to linearized tachyon equation, Yang-Mills equation and so on. The non linear equations are coming out from our consideration only if we restrict the class of boundary interactions even more (or, if short distance behavior depends on a coupling constant, as in the case of having only first derivatives of X in V : $V = \omega^{ij} X^i \partial X^j$). It would be very interesting to find the physical requirements that lead to natural restrictions on boundary interactions. One might think that the reason of linearity of equations of motion is the assumption for operators O being the linear functions of coupling constants: $O = \omega^i O_i$. If we do not impose this condition, our final answer is still given by a one point function of the operator $(\delta - 1)V$:

$$S = \langle (\delta - 1)V \rangle + Z. \quad (19)$$

In the case where we have the subspace of all boundary interactions with only renormalizable (but nonlinear) interactions, our assumptions about the short distance behavior of the exact Green function are completely derivable. This suggests that the question of

linearity is not related to renormalization problems but rather to definition (3). In the lines of the original idea of [1], when the string field theory action is defined by a BV antibracket, what probably has to be done in order to get nonlinear classical dynamics for general boundary interaction, is that the definition of the antibracket has to be modified.

Acknowledgements: I would like to thank E. Verlinde for collaboration in the initial stage of this work and J. Distler, K. Li, V. Periwal and E. Witten for helpful discussions.

References

- [1] E. Witten, Phys.Rev., D46 (1992) 5446.
- [2] E. Witten, Some calculations in background independent off-shell string theory, Princeton Preprint IASSNS-HEP-92/63.
- [3] E. Fradkin and A. Tseytlin, Phys.Lett, B163 (1985) 123.
- [4] A. Abouelsaood, C. G. Callan, C. R. Nappi and S. A. Yost, Nucl. Phys. B280 (1987) 599.
- [5] K. Li and E. Witten, Princeton preprint IASSNS-HEP-93/7.